

MADPH-97-1009

July 1997

THE MUON ELECTRIC DIPOLE MOMENT*

Vernon Barger, Chung Kao

*Department of Physics, University of Wisconsin,
Madison, WI 53706, USA*

Ashok Das

*Department of Physics and Astronomy, University of Rochester,
Rochester, NY 14627, USA*

The electric dipole moment of the muon (d_μ) is evaluated in a two Higgs doublet model with a softly broken discrete symmetry. For $\tan\beta \equiv |v_2|/|v_1| \sim 1$, contributions from two loop diagrams involving the t quark and the W boson dominate; while for $\tan\beta \gtrsim 10$, contributions from two loop diagrams involving the b quark and the τ lepton are dominant. For $8 \gtrsim \tan\beta \gtrsim 4$, significant cancellation occurs among the contributions from two loop diagrams and the one loop contribution dominates for $\tan\beta \sim 7$. For $\tan\beta \gtrsim 15$, the calculated d_μ can be close to the reach of a recently proposed experiment at the Brookhaven National Laboratory.

1 Introduction

For a particle with spin $\vec{s}$, and an electric dipole moment (EDM) $\vec{d} \propto \vec{s}$, the interaction Hamiltonian has the following form:

$$H_I = -\vec{d} \cdot \vec{E}, \quad (1)$$

where $\vec{E}$ is an electric field. This Hamiltonian violates both parity (P) invariance and time-reversal (T) invariance. The invariance of CPT implies that the combined transformation of charge conjugation (C) and parity (P) is not an exact symmetry.^{1,2} The EDM of a fermion (d_f) can be described with an effective Lagrangian

$$\mathcal{L}_{eff} = -\frac{i}{2} d_f \bar{\psi} \sigma_{\mu\nu} \gamma_5 \psi F^{\mu\nu}. \quad (2)$$

In the non-relativistic limit, the Lagrangian becomes $\mathcal{L}_{eff} = -H_I = d_f \vec{s} \cdot \vec{E}$.

In the Standard Model (SM) of electroweak interactions, CP violation is generated by the Kobayashi-Maskawa (KM) phase. The electron EDM³ is about $8 \times 10^{-41} \text{e} \cdot \text{cm}$ and the expected muon EDM is about $2 \times 10^{-38} \text{e} \cdot \text{cm}$. Several experiments have been carried out to search for an electron EDM⁴ (d_e)

*Talk presented by C. Kao at the 2nd International Conference on B Physics and CP Violation, Honolulu, Hawaii, USA, March 1997.

and a muon EDM⁵ (d_μ). At 95% C.L., the experimental upper limits on d_e and d_μ are (i) $|d_e| < 6.2 \times 10^{-27} \text{e} \cdot \text{cm}$, and (ii) $|d_\mu| < 1.1 \times 10^{-18} \text{e} \cdot \text{cm}$.

Recently, a dedicated experiment has been proposed⁶ to measure the muon electric dipole moment at the Brookhaven National Laboratory. This experiment will be able to improve the measurement of the muon EDM by at least 4 orders of magnitude.

2 CP Violation from Higgs Exchange

In multi-Higgs doublet models, a discrete symmetry⁷ is usually required for flavor symmetry to be conserved. In two Higgs doublet models, this discrete symmetry is often chosen to be $\phi_1 \rightarrow -\phi_1$ and $\phi_2 \rightarrow +\phi_2$. If this discrete symmetry is only softly broken⁸, not only can Higgs boson exchange generate CP violation but also flavor changing neutral Higgs interactions can be kept at an acceptable level.

A two Higgs doublet model has doublets ϕ_1 and ϕ_2 . After spontaneous symmetry breaking, there remain five physical ‘Higgs bosons’: a pair of singly charged Higgs bosons ($H^\pm$), two neutral CP-even scalars (H_1 and H_2), and a neutral CP-odd pseudoscalar (A). There are two sources of CP violation in the Higgs potential: (i) the mixing of the A with the H_1 and the H_2 , and (ii) the CP violating interaction of AH^+H^- .

Adopting Weinberg’s parameterization⁹ we can write the following neutral Higgs exchange propagators as

$$\begin{aligned} \langle H_1 A \rangle_q &= \frac{1}{2} \sum_n \frac{\sin 2\beta \text{Im} Z_{0n}}{q^2 - m_n^2} \\ &= \frac{1}{2} \sum_n \left[-\frac{\cos^2 \beta \cot \beta \text{Im} \tilde{Z}_{1n}}{q^2 - m_n^2} + \frac{\sin^2 \beta \tan \beta \text{Im} \tilde{Z}_{2n}}{q^2 - m_n^2} \right] \end{aligned} \quad (3)$$

$$\begin{aligned} \langle H_2 A \rangle_q &= \frac{1}{2} \sum_n \frac{\cos 2\beta \text{Im} Z_{0n} - \text{Im} \tilde{Z}_{0n}}{q^2 - m_n^2} \\ &= \frac{1}{2} \sum_n \frac{\cos^2 \beta \text{Im} \tilde{Z}_{1n} + \sin^2 \beta \text{Im} \tilde{Z}_{2n}}{q^2 - m_n^2} \end{aligned} \quad (4)$$

where the summation is over all the mass eigenstates of neutral Higgs bosons. We will assume that the sums are dominated by the lightest neutral Higgs boson of mass m_0 , and drop the sums and indices. There are relations among the CP violation parameters: $\text{Im} Z_0 + \text{Im} \tilde{Z}_0 = -\cot^2 \beta \text{Im} \tilde{Z}_1$, and $\text{Im} Z_0 - \text{Im} \tilde{Z}_0 = +\tan^2 \beta \text{Im} \tilde{Z}_2$. Employing unitarity constraints, Weinberg has shown that $|\text{Im} Z_1| \leq (1/2) |\tan \beta| (1 + \tan^2 \beta)^{1/2}$ and $|\text{Im} Z_2| \leq (1/2) |\cot \beta| (1 + \cot^2 \beta)^{1/2}$.




Figure 1: Feynman diagrams for leading contributions to the muon EDM.

3 The Muon Electric Dipole Moment

The Feynman diagrams contributing to the muon electric dipole moment are shown in Figure 1. There are many more diagrams involving the W boson that are not shown in this figure. We consider only two-loop diagrams with an intermediate photon since the diagrams involving an intermediate Z boson are highly suppressed by the vector part of the Ze^+e^- couplings.

In a two Higgs doublet model the electron EDM and the muon EDM have contributions from two-loop diagrams containing the top quark (t -loop)¹⁰, the gauge bosons (W -loop)¹¹, and the charged Higgs boson (H -loop)¹².

In the contribution from the charged Higgs boson, CP violation comes from both the neutral Higgs transition propagators and the trilinear vertex AH^+H^- .[†] For $\tan\beta$ larger than about 10, there are dominant contributions from the b quark and the τ lepton¹⁴.

There are several interesting aspects: (i) The contributions from the b and the τ loops are proportional to $(\text{Im } Z_0 + \text{Im } \tilde{Z}_0)$. (ii) For the t -loop, the coefficient of the $\text{Im } \tilde{Z}_0$ is much smaller than that of the $\text{Im } Z_0$. (iii) The W -loop does not contribute to the $\text{Im } \tilde{Z}_0$ term.

The one-loop contribution to the muon EDM (d_μ) is

$$d_\mu^{1\text{-loop}} = \frac{em\sqrt{2}G_F \tan^2\beta}{(4\pi)^2} I(\rho)(\text{Im } Z_0 + \text{Im } \tilde{Z}_0),$$

$$I(\rho) \equiv \rho \int_0^1 dx \frac{x^2}{\rho x^2 - x + 1}, \quad (5)$$

where $\rho = m^2/m_0^2$. In the limit of $m_0 \gg m$ (i.e. $\rho \ll 1$), $I(\rho)$ approaches $-\rho[\ln(\rho) + 3/2]$ and $d_{(e,\mu)}^{1\text{-loop}} \propto (m^3/m_0^2)[\ln(m^2/m_0^2)]$, for $m = m_e, m_\mu$.

[†] Contributions from the $H^\pm$ are not expected to be significant, since recent CLEO measurements on $B(b \rightarrow s\gamma)$ constrain $m_{H^\pm}$ to be at least $3M_W$ in Model II.¹³




Figure 2: The muon EDM for $m_0 = 100$ GeV in units of $\text{Im } Z_0$ and $\text{Im } \tilde{Z}_0$.

Figure 2 shows the muon EDM generated from one-loop diagrams (dash), two-loop diagrams (dash-dot) and their total (solid) in units of (a) $\text{Im } Z_0$ and (b) $\text{Im } \tilde{Z}_0$. Also shown are the contributions from two-loop diagrams involving the W boson (dot), the t quark (dash-dot), the b quark (dash-dot-dot), and the τ lepton (dash-dot-dot-dot), in units of (c) $\text{Im } Z_0$ and (d) $\text{Im } \tilde{Z}_0$. This figure shows the effect of varying $\tan \beta$ on the muon EDM, for the case of $m_0 = 100$ GeV. We note that: (i) for $\tan \beta$ close to one, contributions from two loop diagrams involving the top quark and the W boson dominate; (ii) for $\tan \beta \gtrsim 10$ contributions from two loop diagrams involving the b quark and the τ lepton are dominant; (iii) for $8 \gtrsim \tan \beta \gtrsim 4$, significant cancellation occurs among the contributions from two loop diagrams and one loop diagrams dominate for $\tan \sim 7$; (iv) for $\tan \beta \sim 1$ or $\tan \beta \gtrsim 10$, the muon and the electron EDMs satisfy a simple scaling relation $d_\mu \approx (m_\mu/m_e)d_e$; (v) for $8 \gtrsim \tan \beta \gtrsim 4$, $|d_\mu|$ can be two to three times $|(m_\mu/m_e)d_e|$.

The muon EDM can be expressed as

$$\begin{aligned} d_\mu &= d_A \text{Im} Z_0 + d_B \text{Im} \tilde{Z}_0 \\ &= \frac{d_A + d_B}{2} (\text{Im} Z_0 + \text{Im} \tilde{Z}_0) + \frac{d_A - d_B}{2} (\text{Im} Z_0 - \text{Im} \tilde{Z}_0), \end{aligned} \quad (6)$$

where d_A and d_B are the total coefficients of $\text{Im} Z_0$ and $\text{Im} \tilde{Z}_0$. Applying unitarity constraints we can define the maximal muon EDM¹⁵ ($|d_\mu|_{\text{MAX}}$) as

$$|d_\mu|_{\text{MAX}} = \frac{|d_A + d_B|}{4} \cot \beta (1 + \tan^2 \beta)^{1/2} + \frac{|d_A - d_B|}{4} \tan \beta (1 + \cot^2 \beta)^{1/2}. \quad (7)$$

In Figure 3, we present the maximal value allowed by unitarity for the muon and the electron EDMs, as a function of $\tan \beta$ with $m_0 = 100, 200$ and 400 GeV. Also shown is the experimental upper limit for the electron EDM ($|d_e|_{\text{U.L.}}$).




Figure 3: The maximal EDM allowed by unitarity for (a) the muon and (b) the electron.

In this model, for $\tan\beta \gtrsim 10$, $(m_\mu/m_e)|d_e|_{\text{U.L.}}$ could be treated as an upper limit for the muon EDM since $d_\mu \approx (m_\mu/m_e)d_e$. A measured value of the muon EDM above this bound could arise if the muon EDM and the electron EDM are generated from different sources.

4 Conclusions

Our results may be summarized as follows¹⁵:

1. For $\tan\beta \sim 1$, contributions from two loop diagrams involving the t quark and the W boson dominate; while for $\tan\beta \gtrsim 10$, contributions from two loop diagrams involving the b quark and the τ lepton are dominant.
2. For $\tan\beta \sim 1$ or $\tan\beta \gtrsim 10$, $d_\mu \approx (m_\mu/m_e)d_e$.
3. For $8 \gtrsim \tan\beta \gtrsim 4$, significant cancellations occur among the contributions from two loop diagrams and the one loop contribution dominates for $\tan\beta \sim 7$.
4. For $\tan\beta > 15$ and $m_0 < 300$ GeV, Higgs-boson exchange could produce a muon EDM which is close to the reach of the proposed BNL experiment.

A positive result for the muon EDM measurement could shed light on non-standard CP violation. A negative result, however, could mean

1. that $\tan\beta$ is small than 10; or,
2. the CP violation parameters $\text{Im}Z_0$, and $\text{Im}\tilde{Z}_i$ are smaller than their unitarity bounds; or,
3. the masses of the neutral Higgs scalars and the Higgs pseudoscalar are very close to one another.⁹

Acknowledgments

This research was supported in part by the U.S. Department of Energy under Grants No. DE-FG05-87ER40319 (Rochester) and No. DE-FG02-95ER40896 (Wisconsin), and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

References

1. S.M. Barr and W. Marciano, in *CP Violation*, ed. C. Jarlskog (World Scientific, Singapore, 1989).
2. W. Bernreuther and M. Suzuki, *Rev. Mod. Phys.* **63**, 313 (1991).
3. F. Hoogeveen, *Nucl. Phys. B* **341**, 322 (1990); I.B. Khriplovich and M. Pospelov, *Sov. J. Nucl. Phys.* **53**, 638 (1991); M. Booth, University of Chicago Report, EFI-93-01, (1993).
4. E. Commins *et al.*, *Phys. Rev. A* **50**, 2960 (1994); K. Abdullah *et al.*, *Phys. Rev. Lett.* **65**, 2347 (1990).
5. J. Bailey *et al.*, *J. Phys. G* **4**, 345 (1978); J. Bailey *et al.*, *Nucl. Phys. B* **150**, 1 (1979).
6. Y. Semertzidis *et al.*, E821 Collaboration at BNL, *AGS Expression of Interest: Search for an Electric Dipole Moment of Muon*, September 1996.
7. S.L. Glashow and S. Weinberg, *Phys. Rev. D* **15**, 1958 (1977).
8. G.C. Branco and M.N. Rebelo, *Phys. Lett. B* **160**, 117 (1985); J. Liu and L. Wolfenstein, *Nucl. Phys. B* **289**, 1 (1987).
9. S. Weinberg, *Phys. Rev. D* **42**, 860 (1990).
10. S. Barr and A. Zee, *Phys. Rev. Lett.* **65**, 21 (1990).
11. R.G. Leigh, S. Paban and R.-M. Xu, *Nucl. Phys. B* **352**, 45 (1991); J. Gunion and R. Vega, *Phys. Lett. B* **251**, 157 (1990); D. Chang, W.-Y. Keung and T.-C. Yuan, *Phys. Rev. D* **43**, 14 (1991).
12. C. Kao and R.-M. Xu, *Phys. Lett. B* **296**, 435 (1992).
13. CLEO Collaboration, M.S. Alam *et al.*, *Phys. Rev. Lett.* **74**, 2885 (1995).
14. A. Das and C. Kao, *Phys. Lett. B* **372**, 106 (1996).
15. V. Barger, A. Das and C. Kao, *Phys. Rev. D* **55**, 7099 (1997).